Preliminary Results of Phase II KUNPENG Study of Vebreltinib in Patients (Pts) with Advanced NSCLC Harboring c-MET Alterations

Jin-Ji Yang¹, Yan Zhang², Lin Wu³, Jie Hu⁴, Zhe-Hai Wang⁵, Jing-Hua Chen¹², Jian Fang¹³, Yong Song¹⁴, Wei Zhang¹⁵, Ying Cheng¹⁶, Ren-Hua Guo¹⁷, Xing-Ya Li¹⁸, Anna Li¹, Yi-Long Wu¹ stute, Shandong Provincial Affiliated to Fudan University, Shandong Provincial Oncology, Shandong Cancer Hospital and Institute, Shandong First Medical University, Shandong First Medical University, Chengdu, China; ³Thoracic Medical University, Shandong First Medical University, Shandong Cancer Hospital and Institute, Shandong First Medical University, Shandong Cancer Hospital, China; ⁴Pulmonary and critical care medicine, Zhongshan Hospital Affiliated to Fudan University, Shandong Cancer Hospital, China; ⁴Pulmonary and critical Care medicine, Zhongshan Hospital Affiliated to Fudan University, Shandong First Medical University, Shandong Cancer Hospital, China; ⁴Pulmonary and critical care medicine, Zhongshan Hospital of Sichuan University, Shandong First Medical Unive The spital, 30 Cancer Hospital, 40 Cancer Hospital, 50 Cancer Hospi 14 Lung Cancer Department, Tongji Hospital, University of Education/Beijing, China; 14 Department, Tongji Hospital, University Cancer Department, Tongji Hospital, University of Education/Beijing, China; 14 Department, Tongji Hospital, University of Education/Beijing, China; 14 Department, Peking University Cancer Department, Tongji Hospital, Tongji Hospital, University of Education/Beijing, China; 14 Department, Tongji Hospital, University School of Medicine, Nanjing, China; 14 Department, Tongji Hospital, University of Education/Beijing, China; 14 Department, Tongji Hospital, University Cancer Department, Peking University Cancer Department, Tongji Hospital, University Cancer Department, Peking University Cancer Department, Tongji Hospital, University Cancer Department, Peking University Cancer Department, Tongji Hospital, University Cancer Department, Tongji Hospital, University Cancer Department, Peking University Cancer Department, Tongji Hospital, University Cancer Shanghai Chest Hospital, Shanghai Jiao Tong University, Shanghai, China; ¹⁶Lung Cancer Department, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China: ¹⁷Lung Cancer Department, Jiangsu Provincial People's Hospital Nanjing, China; ¹⁸Oncology Department, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China:

Background


1379P

- MET signaling is implicated in cell proliferation, migration, invasion, and survival^[1]. Genomic alterations in MET can manifest in driving oncogenesis in the form of MET exon 14 skipping mutations (METex14) or other activating mutations, MET gene amplification, gene fusion and MET protein overexpression^[2]. METex14 is an independent prognostic factor associated with poorer survival rates in patients with NSCLC^[3, 4].
- Patients with NSCLC harboring METex14, both previously treated and treatment-naïve, showed objective response rates (ORR) between 49.2% and 68% with existing therapies of MET inhibitors^[5-8]. However, the incidence of treatment-related adverse events (TRAE) of grade 3 or higher was substantial^[5, 6].
- Vebreltinib (bozitinib, APL-101, PLB-1001, CBT-101) is a potent highly selective c-MET inhibitor.
- In a phase I study, Vebreltinib has shown promising results in patients with NGS-identified advanced METex14 skipping NSCLC^[9].
- This phase II study, open-label, multicenter and multi-cohort KUNPENG study evaluate the efficacy and safety of vebreltinib in locally advanced or metastatic NSCLC patients with c-MET alterations (NCT 04258033).

Methods

- Eligible patients were ≥18 years of age, ECOG PS 0~1 with stage IIIB/IV NSCLC.
- Patients in Cohort 1 received 200 mg of Vebreltinib twice daily (28 days as a cycle) until disease progression, death, AE leading to discontinuation or withdrawal of consent.
- Primary endpoint was ORR assessed by blinded independent review committee (BIRC) per RECIST v1.1.
- Secondary endpoints included investigator-assessed (INV) ORR, disease control rate (DCR), duration of response (DoR), time to response (TTR), progression-free survival (PFS) and overall survival (OS).

Figure 1. Study Design



Endpoints					
Primary ORR by BIRC per RECIST v1.1	Safety AE				
Secondary					
ORR by INV					
DCR					
DoR					
TTR					
PFS					
OS		/			

Results


As of 09-Aug-2022, 113 patients were enrolled, among whom 52 patients had METex14 skipping mutations were in Cohort 1. (Table 1.)


Table 1. Baseline characteristics

Variables	Cohort 1 (N=52)	Treatment-naïve (n=35)	Previously-treated (n=17)
Age, years			
Mean (standard deviation)	71.3 (8.3)	71.9 (9.0)	70.0 (6.5)
Median (min, max)	71.0 (51.0, 90.0)	71.0 (53.0, 90.0)	70.0 (57.0, 80.0)
Sex, n (%)			
Male	29 (55.8)	18 (51.4)	11 (64.7)
Female	23 (44.2)	17 (48.6)	6 (35.3)
Ethnicity, n (%)			
Han	49 (94.2)	32 (91.4)	17 (100.0)
Others	3 (5.8)	3 (8.6)	0 (0.0)
ECOG PS, n (%)			
0	5 (9.6%)	5 (14.3)	0 (0.0)
	47 (90.4%)	30 (85.7)	17 (100.0)
Smoking, n (%)			
Current	3 (5.8)	2 (5.9)	1 (5.9)
Former	16 (30.8)	10 (29.4)	6 (35.3)
Never	32 (61.5)	22 (64.7)	10 (58.8)
Histological subtype, n (%)	47 (00 4)		
Adenocarcinoma	47 (90.4)	31(88.6)	16 (94.1)
Squamous carcinoma	1 (1.9)	1 (2.9)	0 (0.0)
Large cell lung cancer	1 (1.9)	0 (0.0)	1 (5.9)
NSCLC, not otherwise specified Staging, n (%)	3 (5.8)	3 (8.6)	0 (0.0)
IIIB	5 (0.6)	5 (11 2)	O(O O)
IIIC	5 (9.6) 4 (7.7)	5 (14.3) 3 (8.6)	0 (0.0) 1 (5.9)
IV	43 (82.7)	27 (77.1)	16 (94.1)
Previous systematic anti-tumor treatment, n (. ,	21 (11.1)	10 (34.1)
Chemotherapy			16 (94.1)
Target therapy			3 (17.6)
Immunotherapy			5 (29.4)
			· (/

ECOG PS, Eastern Cooperative Oncology Group performance status; NSCLC, non-small-cell lung cancer.


• Per BIRC, the primary endpoint ORR was 75% (95% CI: 61.1%~86.0%), with 39 participants achieved CR or PR. The treatment-naïve and previously treated patients displayed an ORR of 77.1% (95% CI, 59.9%~89.6%) and 70.6% (95% CI, 44.0%~89.7%), respectively. (Figure 2.) **Figure 2.** A Treatment exposure and response duration of patients






Per INV, ORR for the FAS population showed a confirmed CR or PR in 69.2% (36/52) of participants. The treatment-naïve and treated populations achieved an ORR of 74.3% (95% CI, 56.7%~87.5%) and 58.8% (95% CI, 32.9%~81.6%), respectively. (Figure

Figure 3. A Treatment exposure and response duration of patients Un-Treated Treated Partial Response Durable responder Response start Response end Continued response -----**A** Each bar represents one subject in the study. A durable responder is a subject who has confirmed response fo at least 183 days (6 months).



Months



Per BIRC, DCR was 96.2% (95% CI, 86.8%~99.5%), median DoR was 15.9 months (95% CI, 9.2~17.8), and median TTR was 1.0 month (95% Cl, 1.0~ 2.8). (Table 2.)


The median PFS was 14.1 months (95% CI, 6.4~17.9) and the median OS was 20.7 months (95% CI, 16.2~NE). (Figure 4.)


Table 2. Efficacy

Variables	BIRC-assessed				Investigator-asse	essed
	Treatment-naïve (n=35)	Previously-treate (n=17)	d All (n=52)	Treatment-naïve (n=35)	Previously-treated (n=17)	d All (n=52)
Objective response rate (%)	27 (77.1)	12 (70.6)	39 (75.0)	26 (74.3)	10 (58.8)	36 (69.2)
95% CI	59.9-89.6	44.0-89.7	61.1-86.0	56.7-87.5	32.9-81.6	54.9-81.3
Best overall response						
Complete response (%)	0	0	0	0	0	0
Partial response (%)*	27 (77.1)	12 (70.6)	39 (75.0)	26 (74.3)	10 (58.8)	36 (69.2)
Stable disease (%)	7 (20.0)	4 (23.5)	11 (21.2)	8 (22.9)	4 (23.5)	12 (23.1)
Progressive disease (%)	1 (2.9)	0	1 (1.9)	1 (2.9)	2 (11.8)	3 (5.8)
Not evaluable (%)	0	1 (5.9)	1 (1.9)	0	1 (5.9)	1 (1.9)
Disease control rate (%)	34 (97.1)	16 (94.1)	50 (96.2)	34 (97.1)	14 (82.4)	48 (92.3)
95% CI	85.1-99.9	71.3-99.9	86.8-99.5	85.1-99.9	56.6-96.2	81.5-97.9
Median duration of response, months (95% CI)	16.5 (9.2-NE)	15.3 (3.7-17.8)	15.9 (9.2-17.8)	16.8 (5.6-19.4)	15.7 (3.7 - NE)	15.7 (7.4-19.4)
Median time to response, months (95% CI)	1.0 (1.0-1.2)	1.9 (0.9-4.5)	1.0 (1.0-2.8)	1.0 (1.0-2.6)	2.7 (0.9-4.4)	1.0 (1.0-2.7)
Progression-free survival						
Median, months (95% CI)	14.5 (6.3-20.3)	7.7 (3.7-20.2)	14.1 (6.4-17.9)	12.0 (6.5-20.3)	8.2 (2.9-NE)	10.5 (6.5-18.2)
6-month progression-free survival (95% CI)	71.4 (53.4-83.5)	70.6 (43.1-86.6)	71.2 (56.8-81.5)	77.1 (59.5-87.9)	64.7 (37.7-82.3)	73.1 (58.8-83.1)
12-month progression-free survival (95% CI)	56.9 (38.9-71.3)	47.1 (23.0-68.0)	53.7 (39.3-66.1)	51.4 (34.0-66.4)	41.2 (18.6-62.6)	48.1 (34.1-60.8)

BIRC, blinded independent review committee; CI, confidence interval; NE, not estimate

Figure 4. Kaplan-Meier estimate of (A) progression-free survival and (B) overall survival.





- Subgroup analysis based on MET amplification, liver metastases, brain metastases and age (Table 3.)
- In patients with brain metastases (n=5), the ORR and DCR were both 100.0% (95% CI, 47.8 to 100.0). The median DoR was 5.6 months (95% CI, 3.7 to NE), and the median PFS was 6.4 months (95% CI, 4.5 to NE).
- Among the six patients with liver metastases, the ORR was 66.7% (95% CI, 22.3 to 95.7), DCR was 100.0% (95% CI, 54.1 to 100.0), and the median DoR was 9.2 months (95% CI, 5.5 to NE). The median PFS was 8.2 months (95% CI, 2.8 to NE), and the median OS was 14.5 months (95% CI, 3.7 to NE).
- Among patients aged 75 years or older (n=21), the ORR was 85.7% (95% CI, 63.7 to 97.0), DCR was 95.2% (95% CI, 76.2 to 99.9), and the median DoR was 19.1 months (95% CI, 9.2 to NE). The median PFS was also 18.2 months (95% CI, 6.3 to NE), and the median OS was not reached (95% CI. 14.4 to NE)
- Patients with co-occurring of MET amplification (n=12, 23.1%) demonstrated an ORR of 100.0% (95% CI, 73.5 to 100.0), a median DoR that was 13.5 months (95% CI, 3.7 to NE), and a median PFS of 14.4 months (95% CI, 4.7 to NE). The ORR was significantly higher in patients with co-occurring of MET amplification than those without (P=0.024), while the PFS was comparable (log-rank P=0.518).

Table 3. Subgroup analysis based on MET amplification, liver metastases, brain metastases and age

Variables	MET amplification + (n=12)	MET amplification - V (n=40)	With liver metastases (n=6)	With brain metastases (n=5)	s ≥75 years (n=21)
Objective response rate (%)	12 (100.0)	27 (67.5)	4 (66.7)	5 (100.0)	18 (85.7)
95% CI	73.5, 100.0	50.9, 81.4	22.3, 95.7	47.8, 100.0	63.7, 97.0
Disease control rate (%)	12 (100.0)	38 (95.0)	6 (100.0)	5 (100.0)	20 (95.2)
95% CI	73.5, 100.0	83.1, 99.4	54.1, 100.0	47.8, 100.0	76.2, 99.9
Median duration of response, months (95% CI)	13.5 (3.7, NE)	15.9 (9.2, 19.1)	9.2 (5.5, NE)	5.6 (3.7, NE)	19.1 (9.2, NE)
6-month progression-free sur- vival rate (95% CI)	66.7 (33.7, 86.0)	73.3 (53.7, 85.7)	75.0 (12.8, 96.1)	40.0 (5.2,75.3)	78.9 (53.2, 91.5)
12-month progression-free	57.1 (25.4, 79.6)	56.5 (37.1, 72.0)	25.0 (0.9, 66.5)	20.0 (0.8, 58.2)	68.4 (42.8, 84.4)
survival rate (95% CI)	14.4 (4.7, NE)	13.1 (6.3, 17.9)	8.2 (2.8, NE)	6.4 (4.5, NE)	18.2 (6.3, NE)
Median progression-free sur- vival, months (95% CI)	NR (13.7, NE)	20.3 (15.2, NE)	14.5 (3.7, NE)	17.9 (4.5, NE)	NR (14.4, NR)

CI, confidence interval, NE, not estimated; NR, not reached.

• The incidence of TRAEs was 77.0% (87/113), with 31.0% (35/113) experiencing TRAEs of grade 3 or above. The most common TRAEs were peripheral edema (56.6%), hypoalbuminemia (22.1%) and hypoproteinemia (19.5%). (Table 4.)

Table 4. Most common treatment-related adverse events ($\geq 20\%$ of patients)

Event, n (%)	Any grade	Grade≥3	Grade 3	Grade 4
At least one TRAE	87 (77.0)	35 (31.0)	32 (28.3)	6 (5.3)
Peripheral edema	64 (56.6)	8 (7.1)	8 (7.1)	0 (0.0)
Hypoalbuminaemia	25 (22.1)	0 (0.0)	0 (0.0)	0 (0.0)
Hypoproteinemia	22 (19.5)	0 (0.0)	0 (0.0)	0 (0.0)

TRAE of any grade considered to be related to study drug by the investigators in at least 10% of patients, or any grade 3 or 4 events reported in any patient are shown. TRAE, treatment-related adverse event. No grade 5 TRAE occurred.

Conclusion

KUNPENG study offers promising evidence of the efficacy and safety of vebreltinib in patients with locally advanced or metastatic ME-Tex14-positive NSCLC. The results highlight capacity of vebreltinib to provide robust and durable responses with a favorable safety profile, signposting its potential as a therapeutic option for METex14-positive NSCLC.

Corresponding author

Yi-Long Wu syylwu@live.cn

Reference

- 1. Smyth EC, F Sclafani, and D Cunningham. Emerging molecular targets in oncology: clinical potential of MET/hepatocyte growth-factor inhibitors. Onco Targets Ther, 2014, 7: 1001-14. DOI: 10.2147/OTT.S44941.
- 2. Drilon A, F Cappuzzo, SI Ou, et al. Targeting MET in Lung Cancer: Will Expectations Finally Be MET? J Thorac Oncol, 2017, 12(1): 15-26. DOI: 10.1016/j.jtho.2016.10.014. 3. Yeung SF, JHM Tong, PPW Law, et al. Profiling of Oncogenic Driver Events in Lung Adenocarcinoma Revealed MET Mutation as Independent Prognostic Factor. J Thorac Oncol,
- 2015, 10(9): 1292-1300. DOI: 10.1097/JTO.000000000000620 4. Tong JH, SF Yeung, AW Chan, et al. MET Amplification and Exon 14 Splice Site Mutation Define Unique Molecular Subgroups of Non-Small Cell Lung Carcinoma with Poor Prognosis.
- Clin Cancer Res, 2016, 22(12): 3048-56. DOI: 10.1158/1078-0432.CCR-15-2061. 5. Yu Y, J Zhou, X Li, et al. Gumarontinib in patients with non-small-cell lung cancer harbouring MET exon 14 skipping mutations: a multicentre, single-arm, open-label, phase 1b/2 trial.
- EClinicalMedicine, 2023, 59: 101952. DOI: 10.1016/j.eclinm.2023.101952. 6 Lu S, J Fang, X Li, et al. Once-daily savolitinib in Chinese patients with pulmonary sarcomatoid carcinomas and other non-small-cell lung cancers harbouring MET exon 14 skipping
- alterations: a multicentre, single-arm, open-label, phase 2 study. Lancet Respir Med, 2021, 9(10): 1154-1164. DOI: 10.1016/s2213-2600(21)00084-9. 7. Paik PK, E Felip, R Veillon, et al. Tepotinib in Non-Small-Cell Lung Cancer with MET Exon 14 Skipping Mutations. N Engl J Med, 2020, 383(10): 931-943. DOI: 10.1056/NEJ-Moa2004407
- 8. Wolf J, T Seto, JY Han, et al. Capmatinib in MET Exon 14-Mutated or MET-Amplified Non-Small-Cell Lung Cancer. N Engl J Med, 2020, 383(10): 944-957. DOI: 10.1056/NEJ-Moa2002787
- 9. Yang J, Q Zhou, H Chen, et al. Abstract CT127: A phase I study of cMET inhibitor bozitinib in patients with advanced NSCLC harboring cMET alterations. Cancer Res, 2020, 80(16 Suppl): CT127. DOI: 10.1158/1538-7445.AM2020-CT127.

DOI# 00010454